

A. Harris
899780

=> fil caplu;e t-lymphocyte/ct 5
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
0.15	0.15

FULL ESTIMATED COST

FILE 'CAPLUS' ENTERED AT 12:14:41 ON 16 AUG 2001
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2001 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications.

FILE COVERS 1947 - 16 Aug 2001 VOL 135 ISS 8
FILE LAST UPDATED: 15 Aug 2001 (20010815/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

This file supports REGISTRY for direct browsing and searching of all substance data from the REGISTRY file. Enter HELP FIRST for more information.

CAPLUS now provides online access to patents and literature covered in CA from 1947 to the present. On April 22, 2001, bibliographic information and abstracts were added for over 2.2 million references published in CA from 1947 to 1966.

The CA Lexicon is now available in the Controlled Term (/CT) field. Enter HELP LEXICON for full details.

Attention, the CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

E#	FREQUENCY	AT	TERM
--	-----	--	----
E1	0	1	T-LYMPHOBLASTIC/CT
E2	0	2	T-LYMPHOBLASTIC LEUKEMIA/CT
E3	0	2 -->	T-LYMPHOCYTE/CT
E4	0	2	T-LYMPHOCYTE (L) HELPER/CT
E5	0	2	T-LYMPHOCYTE CELL GROWTH FACTOR/CT

=> e t-cell/ct 5

E#	FREQUENCY	AT	TERM
--	-----	--	----
E1	0	2	T-CADHERINS/CT
E2	0	15	T-CADINOL/CT
E3	1	2 -->	T-CELL/CT

```

E4          0      2      T-CELL .GAMMA./.DELTA. RECEPTORS/CT
E5         195      3      T-CELL ACTIVATION/CT

```

=> e e3+all

```

E1          1      --> T-cell/CT
E2         19657    USE T cell (lymphocyte)/CT
***** END***

```

=> e leukapheresis/ct 5

```

E#  FREQUENCY  AT  TERM
--  -
E1      1      LEUCTRA TENELLA/CT
E2      0      1  LEUCURUS/CT
E3     29      2 --> LEUKAPHERESIS/CT
E4      0      1  LEUKARTII/CT
E5    22616    48  LEUKEMIA/CT

```

=> s e3+all/ct

THE ESTIMATED SEARCH COST FOR FILE 'CAPLUS' IS 4.44 U.S. DOLLARS
DO YOU WANT TO CONTINUE WITH THIS REQUEST? (Y)/N or END:n
SEARCH ENDED BY USER

=> e e3+all/ct

```

E1      29      --> Leukapheresis/CT
                     HN Valid heading during volumes 86-125 (1977-1996) only.
E2                     NEW Plasmapheresis (L) leukapheresis/CT
***** END***

```

=> s e1-2

```

      29 LEUKAPHERESIS/CT
     519 PLASMAPHERESIS/CT
     102 LEUKAPHERESIS/IT
      55 "PLASMAPHERESIS (L) LEUKAPHERESIS"/CT
L1      84 (LEUKAPHERESIS/CT OR "PLASMAPHERESIS (L) LEUKAPHERESIS"/CT)

```

=> e gm-csf/ct 5

```

E#  FREQUENCY  AT  TERM
--  -
E1      0      2  GM 85 (ZEOLITE)/CT
E2      0      1  GM-CFC/CT
E3      0      2 --> GM-CSF/CT
E4      0      2  GM-CSF RECEPTORS/CT
E5      0      1  GM1/CT

```

=> e granulocyte macrophage colony stimulating factor/ct 5

```

E#  FREQUENCY  AT  TERM
--  -
E1     227     10  GRANULOCYTE COLONY-STIMULATING FACTOR RECEPTORS/CT
E2      0      2  GRANULOCYTE LEUKOCYTE/CT
E3      0      --> GRANULOCYTE MACROPHAGE COLONY STIMULATING FACTOR/CT
E4      0      2  GRANULOCYTE STEM CELL/CT
E5      0      1  GRANULOCYTE-ERYTHROID-MACROPHAGE-MONOCYTE/CT

```

=> e anti-cd3/ct 5

```

E#  FREQUENCY  AT  TERM

```

```

--      -----      --      ----
E1          589      2      ANTI-ALZHEIMER'S DRUGS/CT
E2          1      ANTI-ALZHEIMERS'S AGENTS/CT
E3          0      --> ANTI-CD3/CT
E4          0      2      ANTI-CHEK/CT
E5          0      1      ANTI-GLOMERULAR/CT

```

=> e cd3/ct 5

```

E#      FREQUENCY      AT      TERM
--      -----      --      ----
E1          0      2      CD29 (ANTIGEN)/CT
E2          0      2      CD29 ANTIGENS/CT
E3          0      1 --> CD3/CT
E4         1603      12     CD3 (ANTIGEN)/CT
E5          0      9      CD3 (ANTIGEN) (L) COMPLEXES/CT

```

=> e e4+all/ct

```

E1          0      BT3 Immune factors (non-CA heading)/CT
E2         105281     BT2 Antigens/CT
E3          1769     BT1 CD antigens/CT
E4           477     BT2 Proteins, general/CT
E5         176374     BT1 Proteins, specific or class/CT
E6          1603     --> CD3 (antigen)/CT
                        HN Valid heading during volume 126 (1997) to
                        present.
E7                        OLD Antigens (L) CD3/CT
E8                        UF CD3 antigens/CT
E9                        UF T3 antigen/CT
E10          57      NT1 TCR .alpha..beta.-CD3 complex/CT
E11          203     NT1 TCR-CD3 complex/CT
E12         19657     RT T cell (lymphocyte)/CT
***** END***

```

=> s e6-12

```

1603 "CD3 (ANTIGEN)"/CT
105281 ANTIGENS/CT
4431 CD3/IT
2161 "ANTIGENS (L) CD3"/CT
0 "CD3 ANTIGENS"/CT
0 "T3 ANTIGEN"/CT
57 "TCR .ALPHA..BETA.-CD3 COMPLEX"/CT
203 "TCR-CD3 COMPLEX"/CT
19657 "T CELL (LYMPHOCYTE)"/CT
L2      22701 ("CD3 (ANTIGEN)"/CT OR "ANTIGENS (L) CD3"/CT OR "CD3
ANTIGENS"/C
T OR "T3 ANTIGEN"/CT OR "TCR .ALPHA..BETA.-CD3 COMPLEX"/CT OR
"TCR-CD3 COMPLEX"/CT OR "T CELL (LYMPHOCYTE)"/CT)

```

=> fil reg;e t-lymphocyte/cn 5

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	19.05	19.20

FILE 'REGISTRY' ENTERED AT 12:18:07 ON 16 AUG 2001
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2001 American Chemical Society (ACS)

STRUCTURE FILE UPDATES: 15 AUG 2001 HIGHEST RN 351490-25-0
DICTIONARY FILE UPDATES: 15 AUG 2001 HIGHEST RN 351490-25-0

TSCA INFORMATION NOW CURRENT THROUGH January 11, 2001

Please note that search-term pricing does apply when
conducting SmartSELECT searches.

Structure search limits have been increased. See HELP SLIMIT
for details.

E1 1 T-KOOL 145/CN
E2 1 T-LAK CELL-ORIGINATED PROTEIN KINASE/CN
E3 0 --> T-LYMPHOCYTE/CN
E4 1 T-LYMPHOCYTE SUPPRESSOR FACTOR (HUMAN CLONE .LAMBDA.SUP25
RE DUCED)/CN
E5 1 T-LYMPHOCYTE SUPPRESSOR FACTOR (HUMAN CLONE .LAMBDA.SUP25
RE DUCED), 16-L-ISOLEUCINE-/CN

=> e granulocyte macrophage colony stimulating factor/cn 5

E1 1 GRANULOCYTE COLONY-STIMULATING FACTOR RECEPTOR (HUMAN
SPLICE D 775-AMINO ACID ISOFORM)/CN
E2 1 GRANULOCYTE COLONY-STIMULATING FACTOR RECEPTOR (HUMAN
SPLICE D 873-AMINO ACID ISOFORM)/CN
E3 0 --> GRANULOCYTE MACROPHAGE COLONY STIMULATING FACTOR/CN
E4 1 GRANULOCYTE PEPTIDE A (CATTLE)/CN
E5 1 GRANULOCYTE PEPTIDE A (MOUSE)/CN

=> e "gm-csf"/cn 5

E1 1 GM-30/CN
E2 1 GM-AS/CN
E3 1 --> GM-CSF/CN
E4 1 GM-CSF RECEPTOR (HUMAN .ALPHA.-SUBUNIT SOLUBLE 3)/CN
E5 1 GM-CSF/IL-2 INHIBITION FACTOR (ORF VIRUS STRAIN NZ-2 GENE
GI F)/CN

=> s e3

L3 1 GM-CSF/CN

=> d ide can

L3 ANSWER 1 OF 1 REGISTRY COPYRIGHT 2001 ACS
RN 83869-56-1 REGISTRY
CN Colony-stimulating factor 2 (9CI) (CA INDEX NAME)
OTHER NAMES:
CN Colony-stimulating factor II

CN CSF 2
 CN **GM-CSF**
 CN Granulocyte-macrophage colony-simulating factor
 CN Granulocyte-macrophage colony-stimulating activity
 CN Granulocyte-macrophage colony-stimulating factor
 CN Granulocyte-macrophage-inducing factor
 CN Granulocyte-monocyte colony-stimulating factor
 CN Macrophage-granulocyte CSF
 CN Macrophage-granulocyte-colony-stimulating factor
 MF Unspecified
 CI PMS, MAN
 PCT Manual registration
 LC STN Files: AGRICOLA, AIDSLINE, ANABSTR, BIOBUSINESS, BIOSIS, CA,
 CANCERLIT, CAPLUS, CBNB, CEN, CHEMCATS, CIN, CSCHEM, DRUGPAT,
 DRUGUPDATES, IPA, MEDLINE, MRCK*, PHAR, PROMT, TOXLINE, TOXLIT,
 USPATFULL
 (*File contains numerically searchable property data)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

7918 REFERENCES IN FILE CA (1967 TO DATE)

146 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA

7941 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 135:106281
 REFERENCE 2: 135:106253
 REFERENCE 3: 135:106236
 REFERENCE 4: 135:106161
 REFERENCE 5: 135:106093
 REFERENCE 6: 135:105635
 REFERENCE 7: 135:103358
 REFERENCE 8: 135:103357
 REFERENCE 9: 135:103355
 REFERENCE 10: 135:102853

=> fil medl,caplus,biosis,embase,wpids,jicst

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

6.41

25.61

FILE 'MEDLINE' ENTERED AT 12:20:01 ON 16 AUG 2001

FILE 'CAPLUS' ENTERED AT 12:20:01 ON 16 AUG 2001

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2001 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'BIOSIS' ENTERED AT 12:20:01 ON 16 AUG 2001
COPYRIGHT (C) 2001 BIOSIS(R)

FILE 'EMBASE' ENTERED AT 12:20:01 ON 16 AUG 2001
COPYRIGHT (C) 2001 Elsevier Science B.V. All rights reserved.

FILE 'WPIDS' ENTERED AT 12:20:01 ON 16 AUG 2001
COPYRIGHT (C) 2001 DERWENT INFORMATION LTD

FILE 'JICST-EPLUS' ENTERED AT 12:20:01 ON 16 AUG 2001
COPYRIGHT (C) 2001 Japan Science and Technology Corporation (JST)

=> s (lymphocyte! or cd4(w)cd8(w)ratio or immunity(a)cellular or lymphocyte
cooperation or t cells or thymus dependent lymphocyte or t lymphocyte!)

L4 307125 FILE MEDLINE
L5 128865 FILE CAPLUS
L6 220446 FILE BIOSIS
L7 185921 FILE EMBASE
L8 5634 FILE WPIDS
L9 13998 FILE JICST-EPLUS

TOTAL FOR ALL FILES

L10 861989 (LYMPHOCYTE! OR CD4(W) CD8(W) RATIO OR IMMUNITY(A) CELLULAR OR
LYMPHOCYTE COOPERATION OR T CELLS OR THYMUS DEPENDENT
LYMPHOCYTE
OR T LYMPHOCYTE!)

=> s 12 and 110

L11 5708 FILE MEDLINE
L12 15727 FILE CAPLUS
L13 6 FILE BIOSIS
L14 10 FILE EMBASE
L15 0 FILE WPIDS
L16 0 FILE JICST-EPLUS

TOTAL FOR ALL FILES

L17 21451 L2 AND L10

=> s 117 and 11

L18 9 FILE MEDLINE
L19 3 FILE CAPLUS
L20 0 FILE BIOSIS
L21 0 FILE EMBASE
L22 0 FILE WPIDS
L23 0 FILE JICST-EPLUS

TOTAL FOR ALL FILES

L24 12 L17 AND L1

=> s (l3 or gm csf or granulocyte macrophage colony stimulat?)

L25 13049 FILE MEDLINE
L26 10857 FILE CAPLUS
L27 17420 FILE BIOSIS
L28 15866 FILE EMBASE

L29 759 FILE WPIDS
L30 1428 FILE JICST-EPLUS

TOTAL FOR ALL FILES

L31 59379 (L3 OR GM CSF OR GRANULOCYTE MACROPHAGE COLONY STIMULAT?)

=> s 124 and 131

L32 0 FILE MEDLINE
L33 2 FILE CAPLUS
L34 0 FILE BIOSIS
L35 0 FILE EMBASE
L36 0 FILE WPIDS
L37 0 FILE JICST-EPLUS

TOTAL FOR ALL FILES

L38 2 L24 AND L31

=> d 1-2 cib abs it

'CIB' IS NOT A VALID FORMAT FOR FILE 'CAPLUS'

The following are valid formats:

ABS ----- GI and AB
ALL ----- BIB, AB, IND, RE
APPS ----- AI, PRAI
BIB ----- AN, plus Bibliographic Data and PI table (default)
CAN ----- List of CA abstract numbers without answer numbers
CBIB ----- AN, plus Compressed Bibliographic Data
DALL ----- ALL, delimited (end of each field identified)
DMAX ----- MAX, delimited for post-processing
FAM ----- AN, PI and PRAI in table, plus Patent Family data
FBIB ----- AN, BIB, plus Patent FAM
IND ----- Indexing data
IPC ----- International Patent Classifications
MAX ----- ALL, plus Patent FAM, RE
PATS ----- PI, SO
SAM ----- CC, SX, TI, ST, IT
SCAN ----- CC, SX, TI, ST, IT (random display, no answer numbers;
 SCAN must be entered on the same line as the DISPLAY,
 e.g., D SCAN or DISPLAY SCAN)
STD ----- BIB, IPC, and NCL

IABS ----- ABS, indented with text labels
IALL ----- ALL, indented with text labels
IBIB ----- BIB, indented with text labels
IMAX ----- MAX, indented with text labels
ISTD ----- STD, indented with text labels

OBIB ----- AN, plus Bibliographic Data (original)
OIBIB ----- OBIB, indented with text labels

SBIB ----- BIB, no citations
SIBIB ----- IBIB, no citations

HIT ----- Fields containing hit terms

HITIND ----- IC, ICA, ICI, NCL, CC and index field (ST and IT)
 containing hit terms
 HITRN ----- HIT RN and its text modification
 HITSTR ----- HIT RN, its text modification, its CA index name, and
 its structure diagram
 FHITSTR ----- First HIT RN, its text modification, its CA index name, and
 its structure diagram
 KWIC ----- Hit term plus 20 words on either side
 OCC ----- Number of occurrence of hit term and field in which it occurs

To display a particular field or fields, enter the display field codes. For a list of the display field codes, enter HELP DFIELDS at an arrow prompt (=>). Examples of formats include: TI; TI,AU; BIB,ST; TI,IND; TI,SO. You may specify the format fields in any order and the information will be displayed in the same order as the format specification.

All of the formats (except for SAM, SCAN, HIT, HITIND, HITRN, HITSTR, FHITSTR, KWIC, and OCC) may be used with DISPLAY ACC to view a specified Accession Number.
 ENTER DISPLAY FORMAT (BIB):cbib abs it

L38 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2001 ACS
 2001:265228 Document No. 134:279568 Composition and method of cancer antigen

immunotherapy. Wood, Gary W. (USA). PCT Int. Appl. WO 2001024771 A1 20010412, 50 pp. DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG. (English). CODEN: PIXXD2. APPLICATION: WO 2000-US27399 20001005. PRIORITY: US 1999-412681 19991005.

AB A cancer immunotherapy method and compn. for treating cancer in a patient comprised of vaccinating a patient with a vaccine comprised of the patient's own malignancy and an immunol. adjuvant, removing primed peripheral blood **T lymphocytes** from the patient, stimulating the primed **T lymphocytes** to differentiate into effector **lymphocytes** in vitro, stimulating the effector **T lymphocytes** to proliferate in vitro, and infusing the effector **T lymphocytes** back into the patient. This cancer immunotherapy method can be directed, but is not limited, to the treatment of breast cancer, astrocytoma, and renal cancer.

IT Immunostimulants
 (adjuvants; cancer immunotherapy method comprising a vaccine contg. tumor antigen plus adjuvant, removal of patient's **T cells**, differentiation of **T cells** into effector **T cells**, and infusion of effector **T cells**)

IT Astrocyte
 (astrocytoma, inhibitors; cancer immunotherapy method comprising a vaccine contg. tumor antigen plus adjuvant, removal of patient's **T cells**, differentiation of **T cells**)

- into effector **T cells**, and infusion of effector **T cells**)
- IT Antitumor agents
(astrocytoma; cancer immunotherapy method comprising a vaccine contg. tumor antigen plus adjuvant, removal of patient's **T cells**, differentiation of **T cells** into effector **T cells**, and infusion of effector **T cells**)
- IT Adoptive immunotherapy
Antitumor agents
T cell (lymphocyte)
Vaccines
(cancer immunotherapy method comprising a vaccine contg. tumor antigen plus adjuvant, removal of patient's **T cells**, differentiation of **T cells** into effector **T cells**, and infusion of effector **T cells**)
- IT Interleukin 2
RL: BAC (Biological activity or effector, except adverse); BIOL (Biological study)
(effector **T cell** proliferation induced by interleukin 2)
- IT **T cell (lymphocyte)**
(effector cell; cancer immunotherapy method comprising a vaccine contg. tumor antigen plus adjuvant, removal of patient's **T cells**, differentiation of **T cells** into effector **T cells**, and infusion of effector **T cells**)
- IT Kidney, neoplasm
(inhibitors; cancer immunotherapy method comprising a vaccine contg. tumor antigen plus adjuvant, removal of patient's **T cells**, differentiation of **T cells** into effector **T cells**, and infusion of effector **T cells**)
- IT Antitumor agents
(kidney; cancer immunotherapy method comprising a vaccine contg. tumor antigen plus adjuvant, removal of patient's **T cells**, differentiation of **T cells** into effector **T cells**, and infusion of effector **T cells**)
- IT Plasmapheresis
(leukapheresis; removal of primed **T cells** by)
- IT Antitumor agents
(mammary gland; cancer immunotherapy method comprising a vaccine contg. tumor antigen plus adjuvant, removal of patient's **T cells**, differentiation of **T cells** into effector **T cells**, and infusion of effector **T cells**)
- IT Mammary gland
(neoplasm, inhibitors; cancer immunotherapy method comprising a vaccine contg. tumor antigen plus adjuvant, removal of patient's **T cells**, differentiation of **T cells** into effector **T cells**, and infusion of effector

T cells)
 IT T cell (lymphocyte)
 (proliferation; effector T cell proliferation induced by interleukin
 2)
 IT Antigens
 RL: BAC (Biological activity or effector, except adverse); THU
 (Therapeutic use); BIOL (Biological study); USES (Uses)
 (tumor-assocd.; cancer immunotherapy method comprising a vaccine
 contg.
 tumor antigen plus adjuvant, removal of patient's T
 cells, differentiation of T cells into
 effector T cells, and infusion of effector
 T cells)
 IT Cell differentiation
 (use of anti-CD3 (OKT3) for differentiation of T
 cells into effector T cells)
 IT CD3 (antigen)
 RL: BPR (Biological process); BIOL (Biological study); PROC (Process)
 (use of anti-CD3 (OKT3) for differentiation of T
 cells into effector T cells)
 IT 83869-56-1, Gm-csf
 RL: BAC (Biological activity or effector, except adverse); THU
 (Therapeutic use); BIOL (Biological study); USES (Uses)
 (cancer immunotherapy method comprising a vaccine contg. tumor antigen
 plus adjuvant, removal of patient's T cells,
 differentiation of T cells into effector T
 cells, and infusion of effector T cells)
 IT 140608-64-6, OKT 3
 RL: BAC (Biological activity or effector, except adverse); BIOL
 (Biological study)
 (use of anti-CD3 (OKT3) for differentiation of T
 cells into effector T cells)

L38 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2001 ACS

1998:790654 Document No. 130:13210 Methods and compositions for making
 dendritic cells from expanded populations of monocytes and for activating
 T cells. Nelson, Edward; Strobl, Susan L. (United
 States Dept. of Health and Human Services, USA). PCT Int. Appl. WO
 9853048 A1 19981126, 81 pp. DESIGNATED STATES: W: AL, AM, AT, AU, AZ,
 BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH,
 GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,
 LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,
 SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD,
 RU, TJ, TM; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI,
 FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG.
 (English). CODEN: PIXXD2. APPLICATION: WO 1998-US10311 19980520.
 PRIORITY: US 1997-47348 19970521.

AB Methods of generating IL-3 expanded populations of monocytes and
 differentiating the cells into dendritic cells are provided. Dendritic
 cells are used to activate T cells, in vitro and in
 vivo, and for ex vivo and other therapeutic methods. This facilitates
 the
 use of dendritic cells to generate cell-mediated immune responses.

IT Proteins (specific proteins and subclasses)
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological

process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (BRCA-1; methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

IT Proteins (specific proteins and subclasses)
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (BRCA-2; methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

IT Proteins (specific proteins and subclasses)
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (DCC; methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

IT Proteins (specific proteins and subclasses)
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (FAP; methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

IT Transcription factors
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (FIS (factor for inversion stimulation); methods and compns. for
 making
 dendritic cells from expanded populations of monocytes and for
 activating **T cells**)

IT Proteins (specific proteins and subclasses)
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (GIP; methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

IT Proteins (specific proteins and subclasses)
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (GSP; methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

IT Proteins (specific proteins and subclasses)

RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (HBVc; methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

IT Proteins (specific proteins and subclasses)
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (HBVs; methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

IT Proteins (specific proteins and subclasses)
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (HPV E7; methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

IT Proteins (specific proteins and subclasses)
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (HPV; methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

IT Proteins (specific proteins and subclasses)
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (Hst; methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

IT Antigens
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (Int-2; methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

IT Proteins (specific proteins and subclasses)
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (MAGE-1; methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

IT Proteins (specific proteins and subclasses)
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (MART-1; methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

IT Proteins (specific proteins and subclasses)
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (MEN-1; methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

IT Antigens
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (MUC-1; methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

IT Proteins (specific proteins and subclasses)
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (OB-1; methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

IT Proteins (specific proteins and subclasses)
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (OB-2; methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

IT Proteins (specific proteins and subclasses)
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (RK; methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

IT Proteins (specific proteins and subclasses)
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (ROS; methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

)

IT Proteins (specific proteins and subclasses)
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (TRC; methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

IT Proteins (specific proteins and subclasses)
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (TRP-1 (tyrosinase-related protein 1); methods and compns. for making
 dendritic cells from expanded populations of monocytes and for
 activating **T cells**)

IT Proteins (specific proteins and subclasses)
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (WTI; methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

IT Mycobacterium
 (antigen; methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

IT Carbohydrates, biological studies
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (antigen; methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

IT Proteins (specific proteins and subclasses)
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (cell surface-assocd.; methods and compns. for making dendritic cells
 from expanded populations of monocytes and for activating **T
 cells**)

IT Separation
 (elutriation; methods and compns. for making dendritic cells from
 expanded populations of monocytes and for activating **T
 cells**)

IT Glycophosphoproteins
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (endoplasmic; methods and compns. for making dendritic cells from

expanded populations of monocytes and for activating **T cells**)

IT Proteins (specific proteins and subclasses)
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (gene bcr-c-abl; methods and compns. for making dendritic cells from
 expanded populations of monocytes and for activating **T cells**)

IT RNA formation factors
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (gene myb; methods and compns. for making dendritic cells from
 expanded populations of monocytes and for activating **T cells**)

IT Proteins (specific proteins and subclasses)
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (gene myc; methods and compns. for making dendritic cells from
 expanded populations of monocytes and for activating **T cells**)

IT Lipoproteins
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (gene src; methods and compns. for making dendritic cells from
 expanded populations of monocytes and for activating **T cells**)

IT **Plasmapheresis**
 (leukapheresis; methods and compns. for making dendritic
 cells from expanded populations of monocytes and for activating
T cells)

IT Antitumor agents
 Bacteria (Eubacteria)
 Blood
 Breast tumors
 Cell differentiation
 Colon tumors
 Dendritic cell
 Drugs
 Genetic vectors
 Helper T cell
 Human immunodeficiency virus
 Melanoma
 Monocyte
 Mononuclear cell (leukocyte)

Natural killer cell
 Parasite
T cell (lymphocyte)
 T cell activation
 Tissue culture (animal)
 Transduction (genetic)
 Transformation (genetic)
 Tumors (animal)
 Virus
 (methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)

- IT Antigens
 Antigens
 CD40 ligand
 Carcinoembryonic antigen
 Cytokines
 E6 protein
 Env glycoproteins
 Epidermal growth factor receptors
 Idiotypes (immunoglobulin/TCR)
 Interleukin 1.alpha.
 Interleukin 1.beta.
 Interleukin 3
 Interleukin 4
 Neurofibromin
 Prostate-specific antigen
 Proteins (general), biological studies
 Ras proteins
 Surface antigens
 Tumor necrosis factor .alpha.
 gag proteins
 neu (receptor)
 p53 (protein)
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological
 process); BUU (Biological use, unclassified); BIOL (Biological study);
 PROC (Process); USES (Uses)
 (methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)
- IT Nucleic acids
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL
 (Biological study); PROC (Process)
 (methods and compns. for making dendritic cells from expanded
 populations of monocytes and for activating **T cells**
)
- IT Glial-derived neurotrophic factor
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (receptor, ret; methods and compns. for making dendritic cells from
 expanded populations of monocytes and for activating **T**
 cells)
- IT Neurotrophic factor receptors
 RL: BAC (Biological activity or effector, except adverse); BPR
 (Biological


```

process); BUU (Biological use, unclassified); BIOL (Biological study);
PROC (Process); USES (Uses)
  (ret; methods and compns. for making dendritic cells from expanded
  populations of monocytes and for activating T cells
  )
IT  Epidermal growth factor receptors
    RL: BAC (Biological activity or effector, except adverse); BPR
(Biological
  process); BUU (Biological use, unclassified); BIOL (Biological study);
  PROC (Process); USES (Uses)
    (variant type III; methods and compns. for making dendritic cells from
    expanded populations of monocytes and for activating T
    cells)
IT  9002-10-2, Tyrosinase 83869-56-1, GM-CSF
    RL: BAC (Biological activity or effector, except adverse); BPR
(Biological
  process); BUU (Biological use, unclassified); BIOL (Biological study);
  PROC (Process); USES (Uses)
    (methods and compns. for making dendritic cells from expanded
    populations of monocytes and for activating T cells
    )

```

```

=> s wood g?/au,in
'IN' IS NOT A VALID FIELD CODE
L39      761 FILE MEDLINE
L40      1291 FILE CAPLUS
L41      1080 FILE BIOSIS
'IN' IS NOT A VALID FIELD CODE
L42      615 FILE EMBASE
L43      159 FILE WPIDS
L44      29 FILE JICST-EPLUS

```

```

TOTAL FOR ALL FILES
L45      3935 WOOD G?/AU,IN

```

```

=> s 145 and 117
L46      2 FILE MEDLINE
L47      5 FILE CAPLUS
L48      0 FILE BIOSIS
L49      0 FILE EMBASE
L50      0 FILE WPIDS
L51      0 FILE JICST-EPLUS

```

```

TOTAL FOR ALL FILES
L52      7 L45 AND L17

```

```

=> s 152 not 138
L53      2 FILE MEDLINE
L54      4 FILE CAPLUS
L55      0 FILE BIOSIS
L56      0 FILE EMBASE
L57      0 FILE WPIDS
L58      0 FILE JICST-EPLUS

```

TOTAL FOR ALL FILES
L59 6 L52 NOT L38

=> dup rem l59
PROCESSING COMPLETED FOR L59
L60 6 DUP REM L59 (0 DUPLICATES REMOVED)

=> d cbib abs 1-6

L60 ANSWER 1 OF 6 CAPLUS COPYRIGHT 2001 ACS
2001:175818 Document No. 134:293737 Resistance of Copenhagen rats to
hepatocarcinogenesis does not involve T-cell immunity. **Wood,**
Geoffrey A.; Korkola, James E.; Archer, Michael C. (Department of
Medical Biophysics, University of Toronto, Toronto, ON, M5S 3E2, Can.).
Carcinogenesis, 22(2), 357-359 (English) 2001. CODEN: CRNGDP. ISSN:
0143-3334. Publisher: Oxford University Press.

AB Previously, we have shown that Copenhagen (Cop) rats are highly
resistant,
compared with susceptible F344 rats, to the growth of glutathione
S-transferase 7-7 (GST 7-7) pos. preneoplastic liver lesions following
treatment with a modified resistant hepatocyte (RH) protocol. Donryu
rats, a strain with a level of susceptibility similar to F344, have a
reduced T-cell response compared with the closely related, but highly
resistant, DRH rat. Cop and DRH rats share several characteristics in
their resistance to preneoplastic liver lesion growth and this study,
therefore, was designed to examine whether **T-cells**
play a role in Cop resistance. Cop rats were crossed with an athymic
(nude) rat to produce F1s that were then interbred to produce F2 animals,
some of which were nude with a partial Cop background. A comparison of
the susceptibility of nude F2 animals and their euthymic (non-nude)
littermates allowed us to det. what role, if any, **T-**
cells play in Cop resistance. We treated 11 Cop, 11 F344, 19 nude
F2s, and 18 non-nude F2s with diethylnitrosamine (DEN), followed 3 wk
later by a modified RH protocol. As expected, F344 rats were highly
susceptible, having 41.9 \pm 3.3% (mean \pm SEM) of their liver section
areas occupied by GST 7-7-pos. lesions and Cop rats were highly
resistant,
having only 4.7 \pm 1.1% of their liver section areas occupied by
lesions. Both nude and non-nude F2s were, like Cop rats, highly
resistant
(1.8 \pm 0.29 and 2.7 \pm 0.45%, resp.). These results show that
T-cells are unnecessary for Cop rat resistance, or only
play a minor role, and that the nude parental strain is also likely to be
resistant to the growth of preneoplastic liver lesions.

L60 ANSWER 2 OF 6 CAPLUS COPYRIGHT 2001 ACS
1999:486598 Document No. 131:241295 Target cell range of Haemophilus
ducreyi
hemolysin and its involvement in invasion of human epithelial cells.
Wood, Gwendolyn E.; Dutro, Susan M.; Totten, Patricia A.
(Department of Medicine, Division of Infectious Diseases, University of
Washington, Seattle, WA, USA). Infect. Immun., 67(8), 3740-3749
(English)
1999. CODEN: INFIBR. ISSN: 0019-9567. Publisher: American Society for
Microbiology.

AB Haemophilus ducreyi, the causative agent of chancroid, produces a hemolysin, whose role in virulence is not well defined. To assess the possible role of hemolysin in pathogenesis, we evaluated its target cell range by using wild-type H. ducreyi 35000, nonhemolytic mutants with the hemolysin structural gene deleted, and isogenic strains expressing different amts. of hemolytic activity. The cytotoxicity of the various cell types was assessed by quantitating the release of lactate dehydrogenase into culture supernatants as a measure of cell lysis. In these expts., human foreskin fibroblasts, human foreskin epithelial cells, and, to a lesser extent, HEp-2 cells were lysed by H. ducreyi hemolysin. Hemolysin also lysed human blood mononuclear cells and immune system cell lines including U937 macrophage-like cells, T lymphocytes, and B lymphocytes. In contrast, human polymorphonuclear leukocytes were not sensitive to hemolysin under the conditions tested. We also analyzed the effect of hemolysin on invasion of human epithelial cells and found that H. ducreyi strains expressing cloned hemolysin genes showed a 10-fold increase in invasion compared to the control strain. These data support the hypothesis that the H. ducreyi hemolysin is important in the pathogenesis of chancroid and may contribute to ulcer formation, invasion of epithelial cells, and evasion of the immune response.

L60 ANSWER 3 OF 6 CAPLUS COPYRIGHT 2001 ACS

1999:494086 Document No. 131:156735 Interleukin-12 therapy of cutaneous T-cell lymphoma induces lesion regression and cytotoxic t-cell responses. Rook, Alain H.; Wood, Gary S.; Yoo, Elisa K.; Elenitsas, Rosalie; Kao, David M. F.; Sherman, Matthew L.; Witmer, William K.; Rockwell, Kenneth A.; Shane, Ryan B.; Lessin, Stuart R.; Vonderheid, Eric C. (Department of Dermatology, University of Pennsylvania School of Medicine, Philadelphia, PA, USA). Blood, 94(3), 902-908 (English) 1999. CODEN: BLOOAW. ISSN: 0006-4971. Publisher: W. B. Saunders Co..

AB Progression of cutaneous T-cell lymphoma (CTCL) is assocd. with profound defects in cell-mediated immunity and depressed prodn. of cytokines, which support cell-mediated immunity. Because we have obsd. marked defects in interleukin-12 (IL-12) prodn. in CTCL and because IL-12 is crit. for antitumor cytotoxic T-cell responses, we initiated a phase I dose escalation trial with recombinant human IL-12 (rhIL-12) where patients received either 50, 100, or 300 ng/kg rhIL-12 twice weekly s.c. or intralesionally for up to 24 wk. Ten patients were entered: 5 with extensive plaque, 3 with Sezary syndrome, and 2 with extensive tumors with large cell transformation. One patient with Sezary syndrome dropped out after 1 wk for personal reasons. S.c. dosing resulted in complete responses (CR) in 2 of 5 plaque and partial responses (PR) in 2 of 5 plaque, and 1 of 2 Sezary syndrome (overall response rate CR+PR 5 of 9, 56%). A minor response also occurred in 1 of 5 plaque patients. Intralesional dosing resulted in individual tumor regression in 2 of 2 patients. Biopsy of regressing lesions showed a significant decrease in the d. of the infiltrate in all cases and complete resolu. of the infiltrate among those with clin. lesion resolu. An increase in nos. of CD8-pos. and/or TIA-1-pos. T cells were obsd. on

immunohistochem. anal. of skin biopsy specimens obtained from regressing skin lesions. Adverse effects of rhIL-12 on this regimen were minor and limited and included low-grade fever and headache. One patient discontinued rhIL-12 at week 6 because of depression. These results suggest that rhIL-12 may augment antitumor cytotoxic T-cell responses and may represent a potent and well-tolerated therapeutic agent for CTCL.

L60 ANSWER 4 OF 6 CAPLUS COPYRIGHT 2001 ACS

1997:166252 Document No. 126:237398 Resistance to D5 chemically-induced mammary tumors in Copenhagen x nude-derived F2 athymic rats: evidence that

T-cell immunity is not involved in Copenhagen resistance. Korkola, James E.; Wood, Geoffrey A.; Archer, Michael (Faculty of Med., Univ. of Toronto, Toronto, Can.). Carcinogenesis, 18(1), 53-57 (English) 1997. CODEN: CRNGDP. ISSN: 0143-3334. Publisher: Oxford University Press.

AB Resistance to chem.-induced mammary tumors in the Copenhagen rat is well defined, but the mechanism of resistance has yet to be detd. We have tested whether or not Copenhagen rat resistance is dependent on T-cells, since several lines of evidence supported an involvement of the immune system. We crossed Copenhagen rats with an athymic nude

rat

to produce F1s that were interbred to produce F2 animals, some of which were athymic with partial Copenhagen rat background. A comparison of the mammary tumor incidences between the nude athymic F2 animals and their non-nude littermates allowed us to det. what role, if any, T-cells played in resistance. Following treatment with N-methyl-N-nitrosourea, we obsd. no difference in the tumor incidences between the two groups. Furthermore, the mammary tumor incidences in the F2 nude and non-nude animals was almost zero. These results indicate

that

T-cells are not involved in Cop resistance, and that nude rats are resistant to N-methyl-N-nitrosourea-induced mammary tumorigenesis.

L60 ANSWER 5 OF 6 MEDLINE

91273129 Document Number: 91273129. PubMed ID: 1828937. Most CD8+ cells in skin lesions of CD3+ CD4+ mycosis fungoides are CD3+ T cells that lack CD11b, CD16, CD56, CD57, and human Hanukah factor mRNA. Wood G S; Dubiel C; Mueller C; Abel E A; Hoppe R T; Edinger A; Weissman I; Warnke R A. (Department of Dermatology, Case Western Reserve University, Cleveland, Ohio.) AMERICAN JOURNAL OF PATHOLOGY, (1991 Jun) 138 (6) 1545-52. Journal code: 3RS; 0370502. ISSN: 0002-9440. Pub. country: United States. Language: English.

AB To define further the characteristics of CD8+ cells in skin lesions of CD3+ CD4+ mycosis fungoides (MF), the authors used single- and double-label immunohistologic techniques and in situ hybridization to detect antigens and transcripts associated with certain types of

cytotoxic

or suppressor function. The cytotoxic markers included CD16, CD56, CD57, and an anti-sense probe for human Hanukah factor (HuHf) mRNA. Analysis of 23 cases demonstrated that lesional CD8+ cells were CD3+ T cells that generally lacked expression of any of the cytotoxic markers studied. Analysis of another 10 cases confirmed the CD3+ T-cell lineage of lesional CD8+ cells and demonstrated that these cells also lacked expression of the suppressor-associated marker, CD11b. In

aggregate, these results indicate that most CD8+ cells in CD3+ CD4+ MF skin lesions are of T-cell rather than NK-cell differentiation. Their overall phenotype suggests that they may be major histocompatibility complex (MHC)-restricted cytotoxic **T cells** lacking appreciable levels of HuHF serine protease. Because the induction of CD8+ suppressor **T cells** is mediated by CD4+ **T cells** expressing the CD45RA+ RO- phenotype, CD45 epitope expression was studied in 15 MF cases. The vast majority (13/15)

contained

CD3+ CD4+ tumor cells that were CD45+ RA- RB+ RO+ 2B11+. This phenotype is

consistent with memory **T cells** rather than suppressor-inducer **T cells**, and correlates with the paucity of phenotypically defined suppressor **T cells** in CD3+ CD4+ MF skin lesions.

L60 ANSWER 6 OF 6 MEDLINE

90203337 Document Number: 90203337. PubMed ID: 1690762. Leu-8/CD7 antigen

expression by CD3+ **T cells**: comparative analysis of skin and blood in mycosis fungoides/Sezary syndrome relative to normal blood values. **Wood G S**; Hong S R; Sasaki D T; Abel E A; Hoppe R T; Warnke R A; Morhenn V B. (Department of Dermatology, Stanford University Medical Center, CA.) JOURNAL OF THE AMERICAN ACADEMY OF DERMATOLOGY, (1990 Apr) 22 (4) 602-7. Journal code: HVG; 7907132. ISSN: 0190-9622. Pub. country: United States. Language: English.

AB Deficiencies of Leu-8 and CD7 antigens are exhibited by CD3+ **T cells** in the skin lesions of most patients with mycosis fungoides/Sezary syndrome. To determine whether these antigenic abnormalities are limited to involved skin, we studied Leu-8/CD7 expression in 21 skin lesions of mycosis fungoides/Sezary syndrome obtained from 16 patients and compared them with their peripheral blood leukocytes obtained concurrently. There was no correlation between Leu-8/CD7 values in skin lesions versus blood. Blood values were relatively uniform; most patients had 50% or greater of CD3+, Leu-8+ **T cells** and CD3+, CD7+ **T cells**. In contrast, skin values were highly heterogeneous; most patients lacked expression of Leu-8 or CD7 by the majority of lesional CD3+ **T cells**. Furthermore, Leu-8/CD7 antigen deficiency was present in lesional skin in one patient with mycosis fungoides but not in her concurrently sampled pityriasis lichenoides chronica or blood. These findings suggest that Leu-8/CD7 antigen deficiencies in skin lesions of mycosis fungoides/Sezary syndrome do not represent generalized antigenic abnormalities of CD3+ **T cells** in other body compartments and that within the skin, these deficiencies are disease specific within individual patients with more than one dermatosis. Comparative peripheral blood immunophenotyping of the patients with mycosis fungoides/Sezary syndrome and of the control subjects indicated that the control ranges of CD3+/Leu-8+ and CD3+/CD7+ **T cells** (33% or greater) extend lower than reported previously (60% or greater) and suggested that leukemic involvement in patients with mycosis fungoides/Sezary syndrome may correlate with percentages of CD3+, Leu-8+ and/or CD3+, CD7+ **T cells** that fall below the revised control range.

=> log y

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

60.70

86.31

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE

TOTAL

ENTRY

SESSION

CA SUBSCRIBER PRICE

-3.53

-3.53

STN INTERNATIONAL LOGOFF AT 12:27:04 ON 16 AUG 2001